

*Translation of this copy
does not contain all the claims
as published.*

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

I, JUDITH MARGARET ATKINSON, B.A., M.I.T.I. declare

1. That I am a citizen of the United Kingdom of Great Britain and Northern Ireland, residing at 32 Parkes Way, Blackburn, Lancashire.
2. That I am well acquainted with the French and English languages.
3. That the attached is a true translation into the English language of French Patent Application No. 02 13802 filed on 5th November 2002.
4. That all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that wilful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such wilful false statements may jeopardise the validity of the patent application in the United States of America or any patent issuing thereon.

Declared this 22nd day of December, 2003

J.M. Atkinson.

JUDITH M. ATKINSON

10/533699

JG14 REC'D 03 MAY 2005

FRENCH REPUBLIC
INPI
NATIONAL INSTITUTE FOR INDUSTRIAL PROPERTY

PATENT OF INVENTION
UTILITY CERTIFICATE - CERTIFICATE OF ADDITION

OFFICIAL COPY

The Director General of the National Institute for Industrial Property certifies that the attached document is the true certified copy of an application for an Industrial Property Right filed at the Institute.

Issued in Paris, 21 JUL. 2003

For the Director General of the
National Institute for
Industrial Property
The Head of the Patents Department

(signature)

Martine PLANCHE

NATIONAL INSTITUTE FOR INDUSTRIAL PROPERTY
HEAD OFFICE

26 bis, rue de Saint Petersburg
75800 PARIS Cedex 08
Tel.: 33 (0)1 53 04 53 04
www.inpi.fr

Fax: 33 (0)1 53 04 45 23

National Public Institution created by law
no. 51-444 of 19 April 1951

Deposition of documents DATE 5 NOV. 2002 PLACE 75 INPI PARIS NATIONAL REGISTRATION NO. GIVEN BY THE INPI 0213802 FILING DATE GIVEN BY THE INPI 05 NOV. 2002		1 NAME AND ADDRESS OF THE APPLICANT OR OF THE AUTHORISED AGENT TO WHOM CORRESPONDENCE MUST BE ADDRESSED LES LABORATOIRES SERVIER Patent Department 12, place de La Défense 92415 COURBEVOIE Cedex FRANCE	
Your references for this file (optional) 29049			
Confirmation of a deposit by facsimile <input type="checkbox"/> No. given by INPI to the facsimile			
2 NATURE OF THE APPLICATION		Mark one of the following 4 boxes	
Patent application		<input checked="" type="checkbox"/>	
Application for a Utility Certificate		<input type="checkbox"/>	
Divisional application		<input type="checkbox"/>	
Initial patent application or initial utility certificate application		No.	Date
		No.	Date
Conversion of a European Patent Application Initial patent application		<input type="checkbox"/>	
		No.	Date
3 TITLE OF THE INVENTION (maximum 200 characters or spaces) New imidazopyridine compounds, a process for their preparation and pharmaceutical compositions containing them			
4 DECLARATION OF PRIORITY OR REQUEST FOR THE BENEFIT OF THE FILING DATE OF A PRIOR FRENCH APPLICATION		Country or organisation _____ Date _____ No. _____ Country or organisation _____ Date _____ No. _____ Country or organisation _____ Date _____ No. _____ <input type="checkbox"/> If there are other priorities, mark the box and use the "Continuation" form	
5 APPLICANT		<input type="checkbox"/> If there are other Applicants, mark the box and use the "Continuation" form	
Name or style		LES LABORATOIRES SERVIER	
Forenames			
Legal nature			
SIREN No.			
APE-NAF Code			
Address	Street	12, place de La Défense	
	Postal code and town	92415	COURBEVOIE Cedex
Country		FRANCE	
Nationality		FRENCH	
Telephone no. (optional)		01.55.72.60.00	
Fax no. (optional)		01.55.72.72.13	
E-mail address (optional)			

1st filing

INPI NATIONAL INSTITUTE
FOR INDUSTRIAL PROPERTY

PATENT OF INVENTION
UTILITY CERTIFICATE

REQUEST FOR GRANT 2/2

Reserved for INPI

Deposition of documents DATE 5 NOV. 2002 PLACE 75 INPI PARIS NATIONAL REGISTRATION NO. GIVEN BY THE INPI 0213802			
Your references for this file: <i>(optional)</i>		29049	
6 AUTHORISED AGENT			
Name		RIVIERE	
Forename		François	
Practice or company		LES LABORATOIRES SERVIER	
No. of standing power of attorney and/or of contractual bond			
Address	Street	12, place de La Défense	
	Postal code and town	92415	COURBEVOIE CEDEX
Telephone no. <i>(optional)</i>		01.55.72.60.00	
Fax no. <i>(optional)</i>		01.55.72.72.13	
E-mail address <i>(optional)</i>			
7 INVENTOR(S)			
The inventors are the Applicants		<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No In this case, supply a separate designation of inventorship	
8 SEARCH REPORT		For a patent application only (including division and conversion)	
immediate drawing up or deferred drawing up		<input checked="" type="checkbox"/> <input type="checkbox"/>	
Payment of the fees in instalments		Payment in three instalments, for natural persons only <input type="checkbox"/> Yes <input type="checkbox"/> No	
9 REDUCTION IN FEES		For natural persons only <input type="checkbox"/> Requested for the first time for this invention (attach a notice of non-imposition) <input type="checkbox"/> Requested prior to this deposit (attach a copy of the admissibility decision for this invention or indicate its reference)	
If you have used the "Continuation" form, indicate the number of pages attached			
10 SIGNATURE OF THE APPLICANT OR OF THE AUTHORISED AGENT (Name and position of signatory) François RIVIERE, Patent Agent		STAMP OF THE PREFECTURE OR OF THE INPI (signature)	

Law No. 78-17 of 6 January 1978 relating to information processing, data files and rights applies to the responses made on this form. It guarantees right of access to and correction of the data concerning you at the INPI.

The invention relates to new imidazopyridine compounds, to a process for their preparation and to pharmaceutical compositions containing them.

From the structural point of view, very many examples of imidazopyridine compounds are known in the literature, especially for their therapeutic qualities. By way of example, some compounds are used in the treatment of disorders of the central nervous system (WO 0153263), of viral infections (WO 0100611), of allergies (EP 144101) or of cancers (WO 0244156).

AMP-activated protein kinase (AMPK) is a protein kinase which is involved in the cell response of energy stress. The protein is activated by an increase in the intracellular concentrations of AMP following a fall in the concentration of ATP, during physical exercise, for example.

AMPK phosphorylates and modifies the activity of key enzymes of the carbohydrate metabolism. In fact, AMPK plays an important part in lipogenesis, because it inhibits the synthesis of fatty acids and of cholesterol by inactivating acetyl-CoA carboxylase and HMG coeductase. AMPK reduces the expression of fatty acid synthase (FAS), which controls the synthesis of triglycerides.

In addition, AMPK also reduces the expression of one of the key enzymes of neoglucogenesis (PEPCK), which manifests itself in inhibition of the hepatic production of glucose.


Finally, AMPK increases the utilisation of glucose by facilitating the transport of glucose in the muscle.

All those properties combine to make AMPK a target of choice in the treatment of diabetes and of the metabolic disorders associated therewith, the search for pharmacological activators of AMPK accordingly being of fundamental value to the treatment of those pathologies [see Winder WW and Hardie DG: AMP-activated protein kinase, a metabolic master switch: possible roles in type 2 diabetes; Am. J. Physiol., 40: E1-E10, (1999) and cited references].

The Applicant has now found new imidazopyridine compounds having a novel cycloalkyl-imidazopyridine structure, conferring thereon AMPK-activating properties and, more


precisely, anti-diabetic and anti-hyperlipidaemic properties.

The present invention relates more especially to compounds of formula (I):



wherein:

- 5 • R¹ represents a hydrogen atom, a halogen atom or an alkyl, polyhaloalkyl, cyano, nitro, hydroxycarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl or dialkylaminocarbonyl group,
- R² represents a hydrogen atom, an alkyl group, an optionally substituted aryl group, an optionally substituted heteroaryl group, or a group R²⁰-C(X)- wherein:
 - 10 ➤ R²⁰ represents an alkyl group, an alkoxy group, an amino group, an alkylamino group, a dialkylamino group, an optionally substituted aryl group or an optionally substituted heteroaryl group,
 - X represents an oxygen atom, a sulphur atom, or a group NR²¹ wherein R²¹ represents a hydrogen atom or an alkyl group,
- 15 • R³ represents a hydrogen atom or an alkyl group,
- n represents an integer from 1 to 6 inclusive,



to their enantiomers, diastereoisomers and also to addition salts thereof with a pharmaceutically acceptable acid or base,

it being understood that:


- the term "alkyl" denotes a linear or branched hydrocarbon chain containing from 1 to 6

carbon atoms,

- the term "alkoxy" denotes an alkyl-oxy group in which the alkyl chain, which is linear or branched, contains from 1 to 6 carbon atoms,
- the term "aryl" denotes a phenyl or biphenyl group,
- 5 - the term "polyhaloalkyl" denotes a linear or branched carbon chain containing from 1 to 3 carbon atoms and from 1 to 7 halogen atoms,
- the term "heteroaryl" denotes a group having from 5 to 11 ring members which is monocyclic or bicyclic, in which at least one of the rings is aromatic, and which contains in the monocyclic ring system or in the bicyclic ring system 2 or 3 hetero atoms selected from nitrogen, oxygen and sulphur, and
- 10 - the term "optionally substituted" associated with the expressions aryl and heteroaryl means that the groups in question are substituted by one or two identical or different substituents selected from halogen atoms and the groups alkyl, alkoxy, polyhaloalkyl, hydroxy, cyano, nitro, amino (optionally substituted by one or two alkyl groups) and
- 15 -C(O)R_d wherein R_d represents a group selected from hydroxy, alkoxy and amino, it being understood that the heteroaryl group may also be substituted by an oxo group on the non-aromatic moiety of the heteroaryl.

Among the pharmaceutically acceptable acids there may be mentioned, without implying any limitation, hydrochloric acid, hydrobromic acid, sulphuric acid, phosphonic acid, acetic acid, trifluoroacetic acid, lactic acid, pyruvic acid, malonic acid, succinic acid, glutaric acid, fumaric acid, tartaric acid, maleic acid, citric acid, ascorbic acid, methane-sulphonic acid, camphoric acid etc..

Among the pharmaceutically acceptable bases there may be mentioned, without implying any limitation, sodium hydroxide, potassium hydroxide, triethylamine etc..



An advantageous embodiment of the invention relates to compounds wherein R¹ represents

a hydrogen atom.

Preferred compounds of the invention are those wherein R^2 represents a hydrogen atom or a group $R^{20}-C(O)-$.

5 Another advantageous embodiment relates to compounds of formula (I) wherein R^3 represents a hydrogen atom.


The group R^{20} to which preference is given in accordance with the invention is an alkoxy group and more especially an ethoxy group.

In preferred compounds of the invention, n represents an integer 4, 5 or 6 and more especially 5.

10 An especially advantageous embodiment of the invention relates to compounds of formula (I) wherein R^1 represents a hydrogen atom, R^2 represents a hydrogen atom or a group $R^{20}-C(O)-$ wherein R^{20} represents an alkoxy group, and n is 4 or 5.

Among the preferred compounds of the invention there may be mentioned more especially 3-cycloheptyl-3*H*-imidazo[4,5-*b*]pyridine-2-amine.

15 The invention relates also to a process for the preparation of compounds of formula (I), characterised in that there is used as starting material the compounds of formula (II):




wherein R^1 and n are as defined for formula (I), which compounds of formula (II) are condensed with isothiocyanate compounds (III):




wherein X and R^{20} are as defined for formula (I), to yield the intermediates of

formula (IV):




wherein R^1 , n , X and R^{20} are as defined for formula (I),

5 which compounds of formula (IV) undergo intramolecular cyclisation in a basic medium and in the presence of a suitable catalyst to yield the compounds (I/a):



which are particular cases of the compounds of formula (I) wherein R^1 , n , X and R^{20} are as defined for formula (I),


10 which compounds of formula (I/a) are converted, in an acid medium, into compounds of formula (I/b):



which are particular cases of the compounds of formula (I) wherein R^1 and n are as defined for formula (I),

15 in which compounds of formula (I/b) the amine function can be functionalised in a basic medium, with the aid of an alkyl halide $Alk-Z$ (wherein Alk represents an alkyl group and Z represents a halogen atom), to yield the compounds of formula (I/c):

- 6 -



which are a particular case of the compounds of formula (I) wherein R^1 and n are as defined for formula (I) and Alk is as defined hereinbefore,

5 which compounds of formulae (I/b) and (I/c) may, in a basic medium, optionally in the presence of suitable catalysts, be reacted with $R^2 - Z'$ (wherein R^2 is as defined for formula (I) and Z' represents a nucleofugal group, such as a halogen atom or a trihaloalkyl group) to yield the compounds of formula (I),

which compounds (I/a), (I/b) and (I/c) constitute the totality of the compounds of formula (I) and:

- 10 - which may, where necessary, be purified according to a conventional purification technique,
- which are separated, where necessary, into the stereoisomers according to a conventional separation technique,
- 15 - which are converted, if desired, into their addition salts with a pharmaceutically acceptable acid or base,

it being understood that:

- at any time considered to be appropriate in the course of the process described above, for the requirements of synthesis the carbonyl, amino or alkylamino group(s) of the starting reagents (II) and (III) may be protected and then, after condensation, deprotected,
- 20 - the reagents (II) and (III) are prepared according to known procedures described in the literature.

The compounds exhibit especially an excellent activity in reducing triglyceride and blood glucose levels. Those properties justify their use therapeutically in the treatment and/or prophylaxis of hyperglycaemia, dyslipidaemia and, more especially, in the treatment of

25 non-insulin-dependent, type II diabetes, of obesity, of glucose intolerance and of

complications of diabetes especially in the cardiovascular area.

The activity of those compounds is also recommended for the treatment and/or prophylaxis of other diseases including type I diabetes, hypertriglyceridaemia, metabolic syndrome, insulin resistance, dyslipidaemia in diabetics, hyperlipidaemia and hypercholesterolaemia.

- 5 The present invention relates also to pharmaceutical compositions comprising as active ingredient at least one compound of formula (I), alone or in combination with one or more inert, non-toxic, pharmaceutically acceptable excipients or carriers.

10 Among the pharmaceutical compositions according to the invention there may be mentioned more especially those that are suitable for oral, parenteral and nasal administration, tablets or dragées, sublingual tablets, gelatin capsules, lozenges, suppositories, creams, ointments, dermal gels etc..

15 The useful dose varies according to the age and weight of the patient, the nature and severity of the disorder and the administration route, which may be oral, nasal, rectal or parenteral. Generally, the unit dose ranges from 0.1 to 500 mg per 24 hours, for treatment in from 1 to 3 administrations.

The Examples that follow illustrate the invention, without limiting it in any way. The structures of the compounds described have been confirmed by customary spectroscopic and spectrometric techniques.

20 The starting materials used are known products or are prepared according to known procedures.

PREPARATION 1 : *N*²-Cyclohexyl-2,3-pyridinediamine

Step a : *N*-Cyclohexyl-3-nitro-2-pyridineamine

A mixture composed of 0.1 mol (15.85 g) of 2-chloro-3-nitropyridine and 0.1 mol

(11.50 ml) of cyclohexylamine is heated at 120°C for 4 hours in 250 ml of DMF in the presence of potassium carbonate (13.81 g). The solution is then extracted with 200 ml of ether and the organic phase is washed three times with water. After drying over magnesium sulphate, the ether is evaporated off.

5 Step b : *N*²-Cyclohexyl-2,3-pyridinediamine

800 ml of ethanol, 0.05 mol (11.06 g) of the nitrated compound prepared in the preceding Step and 3.5 g of palladium-on-carbon are introduced into an autoclave. The mixture is heated at 60°C for 30 minutes under 50 kg of hydrogen and then stirred horizontally at ambient temperature for 3 hours in order to homogenise the solution. After that time, the
10 solution is filtered through a Büchner and then a double filter in order to remove the final residues of palladium-on-carbon, and the ethanol is evaporated off under reduced pressure.

PREPARATION 2 : *N*²-Cycloheptyl-2,3-pyridinediamine

The experimental protocol is identical with that used in Preparation 1, starting from cycloheptylamine instead of cyclohexylamine in Step a.

15

EXAMPLE 1 : Ethyl 3-cyclohexyl-3*H*-imidazo[4,5-*b*]pyrid-2-ylcarbamate

Step a : Ethyl {[2-(cyclohexylamino)-3-pyridyl]imino}(diethylamino)methylcarbamate

A mixture of 0.02 mol (3.82 g) of 3-amino-2-cyclohexylaminopyridine from Preparation 1 and 0.02 mol of ethoxycarbonyl isothiocyanate is stirred in 50 ml of DMF at ambient
20 temperature for 3 hours. The solution is then cooled to 0°C, and 0.05 mol of dipropylamine and then 0.02 mol of mercuric chloride are added in succession. After 15 minutes, the ice bath is removed and the solution is stirred at ambient temperature for 4 hours. After diluting the solution with 100 ml of ethyl acetate, filtration is carried out over Celite and the solvents are evaporated off under reduced pressure. The resulting crude product is then
25 recrystallised from acetonitrile.

Step b : Ethyl 3-cyclohexyl-3*H*-imidazo[4,5-*b*]pyrid-2-ylcarbamate

0.0048 mol of the compound prepared in the preceding Step is dissolved in 100 ml of a solution of methanol and 15% sodium hydroxide solution (50/50). After heating the solution at reflux for 3 hours, the methanol is evaporated off. The precipitate so obtained is suction filtered, washed with water and recrystallised from acetonitrile.

Melting point: 264°C

EXAMPLE 2 : Ethyl 3-cycloheptyl-3*H*-imidazo[4,5-*b*]pyrid-2-ylcarbamate

The experimental protocol is identical with that used in Example 1, starting in Step a from 3-amino-2-cycloheptylaminopyridine from Preparation 2 instead of 3-amino-2-cyclohexylaminopyridine.

Melting point: 177°C

EXAMPLE 3 : 3-Cyclohexyl-3*H*-imidazo[4,5-*b*]pyridine-2-amine

The compound of Example 1 is added to 100 ml of a dioxane solution saturated with gaseous HCl, and the solution is heated at reflux for 12 hours. After cooling the solution, the precipitate is suction filtered, washed with sodium bicarbonate and then recrystallised from acetonitrile.

Melting point: 210°C

EXAMPLE 4 : 3-Cycloheptyl-3*H*-imidazo[4,5-*b*]pyridine-2-amine

The experimental protocol is identical with that used in Example 3, using the compound of Example 2 as starting material.

Melting point: 210°C

PHARMACOLOGICAL STUDY**EXAMPLE A : AMPK activity in a cell model : hepatocytes isolated from rats**

5 The hepatocytes are isolated according to the technique of Berry and Friend [J. Cell Biol, 43, 506-520 (1969)]. The AMPK activity was measured according to the method described by Davies *et al.* [Eur. J. Biochem., 186, 123-128 (1989)]. The latter involves phosphorylation starting from [γ -³²P]-ATP of a peptide substrate (SAMS), based on the sequence surrounding the site phosphorylated by the AMPK of the ACC. The reaction for measuring the AMPK activity ends with the deposition of an aliquot of the reaction medium on a phosphocellulose paper (Whatman P81), on which the SAMS peptide is fixed and the radioactivity of which is measured after washing.

10 By way of example, the compound of Example 4 activates AMPK, after 30 minutes at a concentration of 500 μ M, by 312 % (compared with the base value), whereas the same concentration of AICA riboside, used as reference, under the same conditions, activates it by 178 %.

EXAMPLE B : Hypolipaemic activity

20 The products of the invention were tested *in vivo* in the obese ob/ob mouse, used as a model of obesity-associated insulin resistance. By way of example, the compound of Example 4 significantly reduces the triglycerides at 125 mg/kg *per os* whereas, with metformin, the same reduction is obtained at 250 mg/kg *per os*.

In this model, the compounds of the invention have thus been shown to be powerful hypolipaemic agents.

EXAMPLE C : Pharmaceutical composition


Formulation for the preparation of 1000 tablets each containing 5 mg:

25 Compound of Example 45 g

Hydroxypropylmethylcellulose	2 g
Wheat starch	10 g
Lactose	100 g
Magnesium stearate	3 g






CLAIMS

1- Compounds of formula (I):



wherein:

- R^1 represents a hydrogen atom, a halogen atom or an alkyl, polyhaloalkyl, cyano, nitro, hydroxycarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl or dialkylaminocarbonyl group,
- R^2 represents a hydrogen atom, an alkyl group, an optionally substituted aryl group, an optionally substituted heteroaryl group, or a group $R^{20}-C(X)-$ wherein:
 - R^{20} represents an alkyl group, an alkoxy group, an amino group, an alkylamino group, a dialkylamino group, an optionally substituted aryl group or an optionally substituted heteroaryl group,
 - X represents an oxygen atom, a sulphur atom, or a group NR^{21} wherein R^{21} represents a hydrogen atom or an alkyl group,
- R^3 represents a hydrogen atom or an alkyl group,
- n represents an integer from 1 to 6 inclusive,

- the representation R^1 - means: , ,  or ,


their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base,

it being understood that:

- the term "alkyl" denotes a linear or branched hydrocarbon chain containing from 1 to 6 carbon atoms,
- 5 - the term "alkoxy" denotes an alkyl-oxy group in which the alkyl chain, which is linear or branched, contains from 1 to 6 carbon atoms,
- the term "aryl" denotes a phenyl or biphenyl group,
- the term "polyhaloalkyl" denotes a linear or branched carbon chain containing from 1 to 3 carbon atoms and from 1 to 7 halogen atoms,
- 10 - the term "heteroaryl" denotes a group having from 5 to 11 ring members which is monocyclic or bicyclic, in which at least one of the rings is aromatic, and which contains in the monocyclic ring system or in the bicyclic ring system 2 or 3 hetero atoms selected from nitrogen, oxygen and sulphur, and
- the expression "optionally substituted" associated with the expressions aryl and hetero-
15 aryl means that the groups in question are substituted by one or two identical or different substituents selected from halogen atoms and the groups alkyl, alkoxy, polyhaloalkyl, hydroxy, cyano, nitro, amino (optionally substituted by one or two alkyl groups) and -C(O)R_d wherein R_d represents a group selected from hydroxy, alkoxy and amino, it being understood that the heteroaryl group may also be substituted by an oxo group on
20 the non-aromatic moiety of the heteroaryl.

2- Compounds of formula (I) according to claim 1 wherein the representation



corresponds to , their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.

3- Compounds of formula (I) according to either claim 1 or claim 2 wherein R¹ represents
25 a hydrogen atom, their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.

4- Compounds of formula (I) according to any one of claims 1 to 3 wherein R² represents a hydrogen atom, their enantiomers, diastereoisomers and also addition salts thereof with a

pharmaceutically acceptable acid or base.



5- Compounds of formula (I) according to any one of claims 1 to 3 wherein R^2 represents a group $R^{20}-C(O)-$, their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.

5 6- Compounds of formula (I) according to any one of claims 1 to 5 wherein R^3 represents a hydrogen atom, their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.

10 7- Compounds of formula (I) according to any one of claims 1 to 3, 5 or 6 wherein R^{20} represents an alkoxy group, their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.

8- Compounds of formula (I) according to any one of claims 1 to 7 wherein n represents an integer from 4 to 6 inclusive, their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.


9- Compounds of formula (I) according to any one of claims 1 to 8 wherein

15  represents  , R^1 represents a hydrogen atom, R^2 represents a hydrogen atom


or a group $R^{20}-C(O)-$ wherein R^{20} represents an alkoxy group, and n is 4 or 5, their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.

20 10- Compound of formula (I) according to any one of claims 1 to 4, 6, 8 or 9 which is 3-cycloheptyl-3*H*-imidazo[4,5-*b*]pyridine-2-amine.


11- Process for the preparation of compounds of formula (I) according to claim 1, characterised in that there is used as starting material a compound of formula (II):



wherein R^1 and n are as defined for formula (I), which compounds of formula (II) are condensed with isothiocyanate compounds (III):




5 wherein X and R^{20} are as defined for formula (I), to yield the intermediates of formula (IV):




wherein R^1 , n , X and R^{20} are as defined for formula (I),

10 which compounds of formula (IV) undergo intramolecular cyclisation in a basic medium and in the presence of a suitable catalyst to yield the compounds (I/a):




which are particular cases of the compounds of formula (I) wherein R^1 , n , X and R^{20} are as defined for formula (I),

15 which compounds of formula (I/a) are converted, in an acid medium, into compounds of formula (I/b):



which are particular cases of the compounds of formula (I) wherein R^1 and n are as defined for formula (I),

in which compounds of formula (I/b) the amine function can be functionalised in a basic medium, with the aid of an alkyl halide $Alk-Z$ (wherein Alk represents an alkyl group and Z represents a halogen atom), to yield the compounds of formula (I/c):



which are a particular case of the compounds of formula (I) wherein R^1 and n are as defined for formula (I) and Alk is as defined hereinbefore,

which compounds of formulae (I/b) and (I/c) may, in a basic medium, optionally in the presence of suitable catalysts, be reacted with $R^2 - Z'$ (wherein R^2 is as defined for formula (I) and Z' represents a nucleofugal group, such as a halogen atom or a trihaloalkyl group) to yield the compounds of formula (I),

which compounds (I/a), (I/b) and (I/c) constitute the totality of the compounds of formula (I) and:

- which may, where necessary, be purified according to a conventional purification technique,
- which are separated, where necessary, into the stereoisomers according to a conventional separation technique,
- which are converted, if desired, into their addition salts with a pharmaceutically acceptable acid or base,

it being understood that:

- at any time considered to be appropriate in the course of the process described above, for the requirements of synthesis the carbonyl, amino or alkylamino group(s) of the starting reagents (II) and (III) may be protected and then, after condensation, deprotected,
- 5 - the reagents (II) and (III) are prepared according to known procedures described in the literature.

12- Pharmaceutical composition comprising as active ingredient at least one compound according to any one of claims 1 to 10, alone or in combination with one or more inert, non-toxic, pharmaceutically acceptable excipients or carriers.

- 10 13- Pharmaceutical composition comprising as active ingredient at least one compound according to claim 12 comprising at least one active ingredient according to any one of claims 1 to 10 for use in the manufacture of medicaments for use as an AMPK activator.

- 15 14- Pharmaceutical composition according to claim 12 comprising at least one active ingredient according to any one of claims 1 to 10 for use in the manufacture of medicaments that treat non-insulin-dependent, type II diabetes, obesity, type I diabetes, hyperlipidaemia, hypercholesterolaemia and their cardiovascular complications.

15- Pharmaceutical composition according to claim 12 comprising at least one active ingredient according to any one of claims 1 to 10 for use in the manufacture of medicaments that treat type I and II diabetes and its cardiovascular complications.

- 20 16- Pharmaceutical composition according to claim 12 comprising at least one active ingredient according to any one of claims 1 to 10 for use in the manufacture of medicaments that treat type I and II diabetes.